

MICHAŁ M. SKOCZYLAŚ¹, ANETA URUSKA², IWONA WOJTOWICZ², KATARZYNA ŻYWICKA², ALEKSANDRA BRZozowska³, LIDIA PUCHALSKA-NIEDBAŁ⁴, MARCIN SAWICKI¹, ANNA WALECKA¹, JERZY MALINOWSKI

¹Department of Diagnostic Imaging and Interventional Radiology, Autonomous Public Clinical Hospital No. 1, ²Students' Scientific Circle in Department of Diagnostic Imaging and Interventional Radiology Pomeranian Medical University in Szczecin, Unii Lubelskiej 1, 71-252 Szczecin, ³Outpatient Clinic of Dental Surgery, University Dental Clinic, ⁴Department of Ophthalmology, Pomeranian Medical University in Szczecin, Powstańców Wlkp. 72, 70-111 Szczecin, e-mail: sklm@wp.pl

ABSCESSSES OF INTERNAL ORGANS AS A SIGN OF INTERACTIONS BETWEEN HUMAN ORGANISM AND ENVIRONMENT

Summary

Development of abscesses in humans is conditioned by the presence of microorganisms in the living environment, but this is not the only one environmental factor affecting the frequency of their occurrence. Data indicating the importance of atmospheric pressure and temperature for the development of abscesses in walls and structures of the oral cavity was published overpast several dozen years. There are also reports of the incidence of orbital abscesses and abscesses of the liver and spleen depending on the seasons. Taking into account the modifying effect of the environment on the incidence of purulent inflammation broadens the range of possible population studies and clinical thinking.